


# SENSORY CHANGES AFTER BRAIN INJURY



Christina Ball, OTR/L, CBIS

---

---

---

---

---

---

---

---

1

## BACKGROUND

- Bachelor of Arts in Psychology, Bard College
- Master of Science in Occupational Therapy, New York University
- Certified Brain Injury Specialist (CBIS) through the Brain Injury Association of America (BIAA) since 2021
- Experience in inpatient (acute care, ICU) and outpatient rehabilitation, adult and pediatric populations

SENSORY CHANGES AFTER BRAIN INJURY

---

---

---

---

---

---

---

---

2

## DISCLAIMERS

I have no financial or non-financial interests to disclose.

SENSORY CHANGES AFTER BRAIN INJURY

---

---

---

---

---

---

---

---

3

## OBJECTIVES

1. Discuss common sensory changes occurring in individuals following brain injury.
2. Identify effective strategies to manage sensory changes in acute and chronic phases of brain injury.
3. Examine current strategies for effective education of the individual, caregivers, and other medical professionals regarding sensory changes following brain injury.

SENSORY CHANGES AFTER BRAIN INJURY

---

---

---

---

---

---

---

---

4

## I. SENSORY CHANGES AFTER BRAIN INJURY

### Five Senses by Judith Wright

Now my five senses  
gather into a meaning  
all acts, all presences;  
and as a lily gathers  
the elements together,  
in me this dark and shining,  
that stillness and that moving,  
these shapes that spring from  
nothing,  
become a rhythm that dances,  
a pure design.

While I'm in my five senses  
they send me spinning  
all sounds and silences,  
all shape and colour  
as thread for that weaver,  
whose web within me growing  
follows beyond my knowing  
some pattern sprung from nothing-  
a rhythm that dances  
and is not mine.

---

---

---

---

---

---

---

---


5



## SENSORY CHANGES

Sensory deficits following brain injury can include vestibular, auditory, visual, tactile, gustatory, olfactory, and interoceptive challenges.

More than **80%** of patients with acquired brain injury (ABI) have reported sensory stimuli as very overwhelming to experience since their injury (Thielen et al., 2024).



(Faltynek et al., 2019)

Unfortunately, how individuals with brain injury are processing and registering sensory information from their sensory environment is often missed...

SENSORY CHANGES AFTER BRAIN INJURY

---

---

---

---

---

---

---

---

6



## SENSORY CHANGES

Seeing, hearing, feeling is used to understand the world around us. Registering and processing sensory input guides more complex behaviors and actions (movement, thought, memory).

Animal studies suggest disrupted sensorimotor processing in mild-moderate TBI through demonstration of abnormal sensory behavior, impacted speeded motor tasks, and response time tasks (Alwis & Rajan, 2014).

Sensory processing changes following brain injury may not only affect cognitive and motor deficits, but also contribute to increased post-traumatic headache and migraine (Hoffman et al., 2019).

“All this sensory input, which begins in the brain, has its effect throughout the body.”

– Norman Cousins

SENSORY CHANGES AFTER BRAIN INJURY

---

---

---

---

---

---

---

---

7

## MECHANISM OF ACTION



A big question mark...

- Reduced processing speed and increased sensory sensitivity (de Sain et al., 2022)
- Insular lesions, functional changes in brain activity (Preusser et al., 2015)
- Sensory gating impairment (Kumar et al., 2005)
- Disruption in the integration of sensory information (Brosseau-Lachaine et al., 2008)
- Behavioral mechanisms, attention deficits, sensory threshold changes (Thielen et al., 2022)
- Anxiety hypothesis suggesting sympathetic over-arousal leading to hypervigilance to environmental stimuli (de Sain et al., 2023).

SENSORY CHANGES AFTER BRAIN INJURY

---

---

---

---

---

---

---

---

8

## VISION



- Photophobia (light sensitivity)
- Diplopia (double vision)
- Poor depth perception
- Blurriness
- Binocular dysfunction (convergence insufficiency)
- Overstimulation in visually stimulating environments (clutter, patterns, busy places such as the grocery store, mall, or restaurants)
- Intolerance for visual close work (computer screen, reading, writing, etc.)
- Oculomotor and accommodative changes (impaired eye motility, difficulty tracking, focusing on objects at different distances)
- Visual field loss
- Visual acuity loss
- Impaired pupillary function

SENSORY CHANGES AFTER BRAIN INJURY

---

---

---

---

---

---

---

---

9

## VISION



Visual changes or vision loss can occur due to injury to the eye, eye orbit, central visual deficits (CN lesions), or damage to the eye muscles.

Binocular dysfunction may be related to damage to afferent, efferent, or other visual pathways in the brain controlling central vision and are common in over **40%** of the population (Faltynek et al., 2019; Lew et al., 2009).

However, vision loss can also occur without direct injury to the eye...

Visual deficits identified in TBI are usually focused on impaired oculomotor function, acuity, visual field, and binocularity. There is little research on the **perception** of visual stimuli or the **integration** of visual information and impact on function following TBI.

SENSORY CHANGES AFTER BRAIN INJURY

---

---

---

---

---

---

---

---

10

## VISION



Etiologies of photophobia include: ocular, orbital, visual pathology, certain medications, neurological abnormalities, psychiatric disorders (Abusamak & Alrawashdeh, 2021).

In the general population, prevalence of photophobia is estimated at **10%**. Head trauma (concussion, PCS) is associated with visual symptoms in **69-82%** of patients.

- Photophobia is most severe 1-3 weeks after trauma but can persist up to 6 months or years post head trauma. Most patients rate their photosensitivity as severe. Other common symptoms of light sensitivity include asthenopia, squinting, and headaches.

A prospective study by Brosseau-Lachaine, et al. (2008) showed specific perceptual deficits for complex visual information in children with mTBI despite a normal neuro exam at time of hospital discharge. This discrepancy in perceptual performance was maintained throughout the assessed period (up to 3 months post-injury).

SENSORY CHANGES AFTER BRAIN INJURY

---

---

---

---

---

---

---

---

11

## AUDITORY



- Hypersensitivity to certain sounds/frequencies (hyperacusis)
- Phonophobia (persistent, abnormal fear or aversion to sound)
- Misophonia (strong dislike, hatred of certain sounds)
- Auditory processing deficits
  - Difficulty processing oral information
  - Difficulty filtering out background noise
- Tinnitus
- Hearing loss (conductive, sensorineural, mixed)
- *DSI (dual sensory impairment with vision impairment)*

SENSORY CHANGES AFTER BRAIN INJURY

---

---

---

---

---

---

---

---

12

## AUDITORY



Lack of consistent terminology used to describe auditory symptoms associated with head injury adds to difficulty in diagnosing and addressing impairments, and accurate estimation of prevalence of auditory symptoms in concussion patient populations. Taxonomy of auditory symptoms likely a combination of peripheral, subcortical, and cortical maladaptive changes that result in persistence of auditory symptoms (Theodoroff et al., 2022).

Auditory deficits themselves can arise from 2 main mechanisms: Injury to area of the brain responsible for auditory processing (central), and injury to the physical structure of the ear or inner ear (peripheral) (Faltynek et al., 2019).

SENSORY CHANGES AFTER BRAIN INJURY

---

---

---

---

---

---

---

---

13

## AUDITORY



Studies show that injuries to the auditory system can account for about **20%** of the injuries reported in troops. Studies also note hearing loss in about **70%** of those who experienced a blast and were referred for audiologic evaluation (Lew et al., 2010).

Lew et al. (2007) showed that **38%** of patients with blast-related TBI reported tinnitus compared to non-blast related TBI (**18%**). **62%** of patients with blast-related TBI also reported hearing loss ( $p=0.04$ ).

Noise sensitivity in the general population is approximately **15%** in adults. **35%** of mTBI patients 3 months post injury report noise sensitivity. Noise sensitivity has been found to be a strong predictor of prolonged post-concussive symptoms and has been linked to longer treatment and rehabilitation times post-TBI (Shepherd et al., 2019).

SENSORY CHANGES AFTER BRAIN INJURY

---

---

---

---

---

---

---

---

14

## VESTIBULAR & PROPRIOCEPTION



- Dizziness
  - Disequilibrium
  - Vertigo
  - Lightheadedness
  - Presyncope
- Imbalance (unsteadiness)
- Clumsiness (incoordination)
- Poor postural control and body awareness
- Poor tolerance for positional changes
- Motion sickness

SENSORY CHANGES AFTER BRAIN INJURY

---

---

---

---

---

---

---

---

15

## VESTIBULAR & PROPRIOCEPTION



Benign paroxysmal positional vertigo (BPPV) is the most common cause of dizziness following TBI; however, vestibular diagnoses are of unknown origin in **25%** of chronic TBI cases.

Temporal bone fractures are another injury which can cause significant damage to the inner ear and can cause labyrinthine concussion (presence of hearing loss and vertigo). Another major type of peripheral complication resulting in vestibular dysfunction is perilymphatic fistula.

Central nervous system causes of vertigo and dizziness can stem from post-traumatic seizures, anxiety disorders, or damage to the central vestibular structures (cerebellum, brainstem, or vestibular nuclei).

SENSORY CHANGES AFTER BRAIN INJURY

(Faltynek et al., 2019)

---

---

---

---

---

---

---

---

16

## VESTIBULAR & PROPRIOCEPTION



Balance difficulties following brain injury can have a peripheral vestibular or central nervous system origin. The research literature shows **23-81%** of individuals report balance issues following ABI (Faltynek et al., 2019).

Balance impairment can be attributed to proprioceptive deficits and reduced muscle tone; approximately **65%** of patients with stroke experience loss of proprioception, tactile sensation, and protective reaction (Lee et al., 2015).

Dizziness can present with several different characteristics and result from deficit from the inner ear, brain, cervical spine, or integration of afferent input and tuning with the sensorimotor control system (Hammerle et al., 2019).

SENSORY CHANGES AFTER BRAIN INJURY

---

---

---

---

---

---

---

---

17

## VESTIBULAR & PROPRIOCEPTION



Chamelian & Feinstein (2004):

- 6-month cross-sectional study of 207 adults with mild-moderate TBI, **66.7%** had subjective complaint of post-traumatic dizziness.
- Dizziness was closely linked to psychological distress at 6 months following injury.
- Dizziness emerged as an independent predictor for failure to return to work with rates at 6 months reduced from **75%** (in those without vestibular dysfunction) to **33%**.
- Psychosocial functioning (GHQ, RHFUQ, RTW) was significantly worse in those with dizziness ( $p < .01$ ).

SENSORY CHANGES AFTER BRAIN INJURY

---

---

---

---

---

---

---

---

18

## OLFACTORY & GUSTATORY



- Hypersensitivity to certain smells, fragrances (hyperosmia)
- Dysosmia (distorted odor perception)
- Phantosmia/Parosmia ('false' smells, perceiving smells that aren't there)
- Hyposmia (partial loss or decreased sense of smell)
- Anosmia (loss of smell)
- Hypersensitivity to certain tastes
- Dysgeusia (distortion or decreased sense of taste)
- Ageusia (loss of taste)
- Loss of appetite

SENSORY CHANGES AFTER BRAIN INJURY

---

---

---

---

---

---

---

---

19

## OLFACTORY & GUSTATORY



Taste and smell are the 2 primary chemosensory systems. Patients with altered taste and smell perception after BI are at risk for malnutrition, associated complications (gastrointestinal bleeding, pneumonia, other infection), and poorer outcomes impacting rehabilitation and quality of life (Dutta et al., 2013; Proskynitopoulos, 2016).

Impaired taste perception can result from lesions in the pons, specific thalamic nuclei, and insular cortices. Studies suggest central lesions cause disorders of taste perception whereas peripheral nerve lesions cause altered taste interpretation (Dutta et al., 2013).

Pathophysiological mechanisms cited in OD include peripheral lesions involving olfactory fibers arising from the cribriform plate, impaired cranial relay stations, central pathway involvement...brain regions frequently associated with OD include the olfactory bulb, olfactory tract, temporal lobe, and sub-frontal lobe (Proskynitopoulos, et al., 2016).

SENSORY CHANGES AFTER BRAIN INJURY

---

---

---

---

---

---

---

---

20

## OLFACTORY & GUSTATORY



Patients affected by smell and taste dysfunction report decreased quality of life:

- Changes in food preferences
- Find certain foods less palatable
- Changes in dietary habits
- Unintentional weight loss
- Adverse health effects secondary to changes in diet (hypertension, malnutrition, etc.)
- Decreased energy level (fatigue)
- Avoidance of social activities
- Depressive symptoms

Impaired smell occurs more frequently in patients diagnosed with major depression and associated with higher scores on depression inventories (Proskynitopoulos, et al., 2016).

SENSORY CHANGES AFTER BRAIN INJURY

---

---

---

---

---

---

---

---

21

## TACTILE



- Hypersensitivity to touch
- Paresthesia (pins and needles sensation)
- Pain
- Hypoesthesia (numbness)
- Increased or decreased response to temperature
- Decreased touch discrimination
- Impaired stereognosis

SENSORY CHANGES AFTER BRAIN INJURY

---

---

---

---

---

---

---

---

22

## TACTILE



Impairment in sensory perception can commonly occur following stroke resulting in the experience of negative symptoms (numbness or lack of sensation, inability to determine limb position in space, recognize a held object) and positive symptoms (paresthesia, pins/needles, pain).

This can have a serious impact on all levels of motor production (planning, correction, execution), which involve the use of sensory feedback from the environment.

Reviewing the literature, changes in proprioception and touch occurred in **25-85%** of patients immediately following stroke, and proprioception (**34-63%**) and stereognosis (**31-89%**) modalities were more commonly affected than touch (**7-53%**).

SENSORY CHANGES AFTER BRAIN INJURY

(Gandhi et al., 2020)

---

---

---

---

---

---

---

---

23

## TACTILE



Chronic pain is a common complication of TBI, contributing to morbidity and poor recovery following injury. Mild TBI is associated with a **75.3%** pain rate, moderate-severe TBI associated with a **32.1%** pain rate (Nampiarampil, 2008).

Little research concerning the mechanisms supporting pain and evidence-based guidelines for treating TBI-related pain. Potential mechanisms include neuroinflammation, excitotoxicity, and axonal degeneration (Irvine & Clark, 2018).

In an animal study by Wattiez et al. (2021), tactile hypersensitivity in cephalic and extra-cephalic areas develops early, persisting for at least 9 weeks in multimodal TBI; in closed head TBI, cephalic tactile hypersensitivity is transient lasting 1-7 days after injury. Following, mice display prolonged tactile hypersensitivity in response to normally sub-threshold doses of migraine triggers.

SENSORY CHANGES AFTER BRAIN INJURY

---

---

---

---

---

---

---

---

24



## INTEROCEPTION



- Poor thermoregulation
- Changes in thirst, hunger (appetite)
- Increased pain sensitivity
- Difficulty differentiating between physical sensations (pain vs. itch)
- Heart rate changes
- Difficulty recognizing level of physical exertion
- Difficulty with self-regulation (emotional-regulation)

SENSORY CHANGES AFTER BRAIN INJURY

---

---

---

---

---

---

---

---

25

## INTEROCEPTION



In a study by Pistoia et al. (2015):

- Processing, representation, and perception of bodily signals (interoception) plays an important role in the subjective experiencing of emotional stress.
- Individuals with SCI (n=10) did not differ from healthy controls (n=20) in recognizing emotions via facial expressions; however, they were significantly less accurate than healthy controls in identifying and judging emotionally evocative scenes eliciting fear and anger.
- Results support the idea that afferent sensory feedback is involved in emotional experience. The higher the level of injury of spinal cord, the potentially greater degree of experienced emotional dysfunction.

SENSORY CHANGES AFTER BRAIN INJURY

---

---

---

---

---

---

---

---

26

## INTEROCEPTION



*Maladaptive interoceptive sense:* Hypervigilance of body sensations, catastrophization.

*Adaptive interoceptive sense:* Attention to, regulation, acceptance, of body sensations.

Desdentado et al. (2023) found individuals with ABI were less likely to ignore unpleasant sensations, have increased severity of depressive symptoms, have greater difficulty in emotional regulation, and exhibit worse performance on emotional awareness than neurotypical individuals. Results indicate benefit in addressing both interoceptive and emotional difficulties in treatments targeting psychosocial sequelae of ABI.

Kharrazian (2015) discusses disruption of the brain-gut axis and alterations in autonomic function in BI; dysautonomia has been found in **8-33%** of individuals with TBI per literature review. A common presentation of dysautonomia is persistent sympathetic overactivity in response to nociceptive stimuli and development of chronic pain.

SENSORY CHANGES AFTER BRAIN INJURY

---

---

---

---

---


---

---

---

27

## II. IDENTIFYING & MANAGING SENSORY CHANGES IN ACUTE AND CHRONIC PHASES OF BRAIN INJURY



---

---

---

---

---

---

---

28



### ACUTE PHASE

The primary goal of acute care (hospitalization) is to keep the patient alive and minimize mortality. Rehabilitative goals center on preparing for the next level of care and longer-term considerations.

Managing cognitive-behavioral issues in acute rehabilitation involves teasing out what the problems are and how to address them (behavioral management, environmental modifications, and pharmacological intervention).

Rehabilitative therapies can play a role in addressing visual dysfunction, vestibular dysfunction, and other somatosensory deficits.

SENSORY CHANGES AFTER BRAIN INJURY (Zasler et al., 2013, p. 388)

---

---

---

---

---

---

---

29

### ENVIRONMENTAL MODIFICATIONS



Non-pharmacological intervention for agitation is typically recommended as a first-line approach (environmental modification, behavioral strategies, education for staff, family).

In a study by Carrier et al. (2021): **99%** (n=305) of clinicians reported using at least 1 non-pharmacological approach to manage agitation with environmental modifications being most common (**98%**), followed by patient program modifications (**94%**), behavior modification techniques (**88%**).

Behavioral strategies and non-pharmacological interventions that target specific triggers of agitation and how it manifests, are important for the patient's rehabilitative progress. Aim is not to just resolve behavioral issues, but to understand underlying causes and equip the individual with appropriate tools to promote participation and re-integration (Janzen et al., 2014).

SENSORY CHANGES AFTER BRAIN INJURY

---

---

---

---

---

---

---

30

## ENVIRONMENTAL MODIFICATIONS

- Reducing stimuli (light, noise, distractions)
  - Placing patient in bed
  - Drawing curtains, turning off lights, dimming brightness of computer monitors
  - Turning off TV/radio
  - Limiting number of visitors
  - Speaking in low volume, slowly
  - Avoid sudden grabbing or touching of the patient, approach from the front
- Obtaining hearing devices, vision aids from home to improve orientation
- Create a familiar environment (allow family to bring in personal possessions)
- Minimize tubes and lines (may cover them – abdominal binder)
- Consistent schedule and staff (timed toileting, sleep hygiene)
- Avoid/minimize restraints (1:1 staff supervision) (Carrier et al., 2021; Janzen et al., 2014)

SENSORY CHANGES AFTER BRAIN INJURY

(Carrier et al., 2021; Janzen et al., 2014)

---

---

---

---

---

---

---

---

31

## ACUTE PHASE (NICU)

"The **Supporting and Enhancing NICU Sensory Experiences (SENSE)** program was developed to promote consistent, age-appropriate, responsive, and evidence-based positive sensory exposures for the preterm infant every day of NICU hospitalization...The mismatch between the underdeveloped coping skills of the infant and the intensely stimulating NICU environment may cause physiologic instability, adversely affect growth and development, and ultimately impact long term neurodevelopmental outcomes [3–7]" (Pineda et al., 2019).

- Combines need for parent engagement in the NICU environment with the infant's need for consistent positive sensory exposures.
- Week-by-week sensory exposure plan outlines what type and timing of exposures are developmentally appropriate.
- Weekly assessments performed by the SENSE administrator or neonatal clinician (nurse, PT, OT, SLP), with log sheets used to track daily, positive sensory exposures.

SENSORY CHANGES AFTER BRAIN INJURY

(Pineda et al., 2019)

---

---

---

---

---

---

---

---

32



## CHRONIC/SUBACUTE PHASE

Patients with post-injury hypersensitivity to sensory stimuli are at risk of:

- Mental health difficulties
  - Sensory processing changes after TBI may contribute to PTSD comorbidity, post-traumatic headache, and migraine (Hoffman et al., 2019).
- Physical health difficulties
- Poorer functional outcomes

Systematic literature review showed a limited number of studies (36%) used a validated questionnaire to assess sensory changes after ABI (Thielen et al., 2024).

- Rivermead Post-concussion Symptom Questionnaire (RPQ) – measures light, noise hypersensitivity, other common post-concussion symptoms
- Sensory Sensitivity Questionnaire (SSQ), Sensory Profile (adult/adolescent, child) (SP), Glasgow Sensory Questionnaire (GSQ) – **not standardized to ABI population**

SENSORY CHANGES AFTER BRAIN INJURY

---

---

---

---

---

---

---

---

33


# CHRONIC/SUBACUTE PHASE



Thielen et al. (2024):

- Post-injury hypersensitivity was reported by **76%** of stroke patients, **89%** of TBI patients, and **82%** of brain tumor patients. Complaints occurred across all modalities (multi-sensory, visual, auditory hypersensitivity most prevalent).
- Study on 818 neuro-typical adults, 341 with ABI showed that **79%** of ABI patients reported experiencing an increase in sensitivity to at least 1 sensory modality that was still present in the month previous to study participation.

**Multi-Modal Evaluation of Sensory Sensitivity (MESSY-NL):** Patient-friendly questionnaire to assess sensory sensitivities across multiple sensory modalities (visual, auditory, tactile, olfactory, gustatory, motion sensitivity, temperature sensitivity).




---

---

---

---

---

---

---

---

---

---

34

# CHRONIC/SUBACUTE PHASE



Cognitive Behavioral Therapy (CBT) is considered a diverse set of problem-specific interventions incorporating physical, psychological, and behavioral approaches to managing pain. The individual is taught to use self-regulation, self-control, and coping strategies for pain, depression, and anxiety associated with chronic headaches.

Other non-pharmacological interventions have been explored to reduce pain and headaches following ABI:

- |                     |             |                      |
|---------------------|-------------|----------------------|
| Biofeedback         | Yoga        | Red light therapy    |
| Relaxation training | Mindfulness | Cryotherapy          |
| Acupuncture         | Meditation  | Craniosacral therapy |

Limited evidence for these interventions to support their efficacy with ABI population. Use may be appropriate on a specific case by case basis.

**SENSORY CHANGES AFTER BRAIN INJURY** (Faltynek et al., 2019)

---

---

---

---

---

---

---

---

---

---

35

# VISION



Due to the variety of visual deficits following ABI, identifying specific deficits and visual intervention is important for positive outcome by an appropriate clinician (ophthalmologist/optometrist, neuro-ophthalmologist, rehabilitative clinicians with expertise in visual rehabilitation).

- Prisms to correct double vision, eye misalignment, other visual processing problems
- Patching to correct double vision
- Vision therapy to improve oculomotor function and visual processing
- Low vision aids and/or training to correct for severe acuity loss, field loss, neglect, or scanning issues

(Faltynek et al., 2019; Lew et al., 2010)

**SENSORY CHANGES AFTER BRAIN INJURY**

---

---

---

---

---

---

---

---

---

---

36

## VISION



Treatment approaches for photophobia include:

- Optical filters, tinted lenses (FL-41 lenses, yellow tinted driving lenses)
- Polarized sunglasses (wear of sunglasses indoors is discouraged)
- Hats, hoodies, other headwear to reduce effects from overhead lighting
- Anti-glare covers for electronic devices
- Non-liquid crystal displays
- Topical ocular lubricants and other medications for neuropathic pain, photophobia (treatment of dry eye, anti-epileptics, serotonin-norepinephrine reuptake inhibitors, and tricyclic antidepressants)
- Sympathetic blocks of the superior cervical ganglion has shown significant results in reducing ocular pain and photophobia in refractory cases

SENSORY CHANGES AFTER BRAIN INJURY (Abusamak & Alrawashdeh, 2022)

---

---

---

---

---

---

---

---

37

## AUDITORY



Referral to audiology is recommended to assess auditory dysfunction potential structural and functional changes that occur in the peripheral and central auditory system post head injury.

Sensorineural and conductive hearing loss will often benefit from use of hearing aid/assistive listening devices (ALD's).

Treatment and management of tinnitus is emerging: Sound therapy, masking relief therapy, hearing aid, relaxation and stress management (CBT), desensitization techniques (hyperacusis), TRT (tinnitus retraining therapy; auditory habituation).

- Important to note the type of tinnitus (subjective, neurologic, somatic, and objective) as treatment may differ.

SENSORY CHANGES AFTER BRAIN INJURY (Faltynek et al., 2019)

---

---

---

---

---

---

---

---

38

## AUDITORY



Auditory training is shown to be an effective method to improve auditory processing. Auditory exercises are provided to the patient that use central processes known to influence listening comprehension in more challenging listening environments.

- A variety of auditory training options are available including software (Earobics, FastForWord, Posit Science Brain Fitness) the patient can use at home and others administered by an audiologist or speech-language pathologist (Dichotic Interaural Intensity Difference training) (Lew et al., 2010).

Hearing protection when exposed to loud sounds (concerts, band practice, etc.) is recommended for patients with noise sensitivity. It is generally not recommended to use hearing protection in quiet environments as this can result in the auditory system becoming more sensitive (Theodoroff et al., 2022).

SENSORY CHANGES AFTER BRAIN INJURY

---

---

---

---

---

---

---

---

39

## VESTIBULAR & PROPRIOCEPTION



Vestibular rehabilitation therapy (VRT) with physical and occupational therapy is the most common management for vestibular dysfunction following brain injury.

Type	Description
Habituation	Habituation exercises operate under the principle that the body will eventually reduce its response to unpleasant or noxious stimuli through exposure.
Adaptation	Encouraging long-term plastic change is part of the adaptive strategy, exercises are designed to reinforce a reorganization of the vestibular system through similar means of exposure.
Substitution	This approach involves using different eye-movement systems to substitute the eye-movement strategies normally used on specific tasks (such as saccade versus smooth eye movement strategies).
Treatment of BPPV	As BPPV is thought to be caused by the inappropriate movement of otoconia (bio-crystals within the semicircular canals). Treatment for BPPV involves a variety of maneuvers attempting to make the crystals responsive to gravity again (e.g., Canalith Repositioning Maneuver).

SENSORY CHANGES AFTER BRAIN INJURY (Faltynek et al., 2019)

---

---

---

---

---

---

---

---

40

## VESTIBULAR & PROPRIOCEPTION



In a study by Hammerle et al. (2019):

- Research shows that dizziness following mTBI improves with sensorimotor control exercises combined with manual therapy for the cervical spine due to abnormal cervical proprioception or afferent input to the sensorimotor control system.
- Retrospective records review of patients treated for dizziness following mTBI showed that patients who received cervical spine proprioception retraining (CSPR) were **30x** more likely to report improvement in dizziness symptoms compared to those who received VRT.
- **85%** of individuals in the CSPR group and only **18%** in the usual care group had dizziness improved, suggesting that by identifying altered CSP with no clear central or peripheral signs present, that CSPR may be beneficial for individuals with dizziness following TBI.

SENSORY CHANGES AFTER BRAIN INJURY

---

---

---

---

---

---

---

---

41

## VESTIBULAR & PROPRIOCEPTION



Study by Lee et al. (2015) of 36 patients post-stroke to examine the effects of exercise image training in treating balance and proprioceptive deficit; 18 patients received proprioceptive training, 18 patients received additional motor imagery training.

- Significant improvements in outcome measures with time ( $p < 0.05$ ), the motor imagery training group had significantly greater improvements than the proprioception training group ( $p < 0.05$ ).

Randomized controlled pilot study by Lim et al. (2019) randomly assigned 30 patients post-stroke to a multi-sensorimotor training group (Reha-Bar, balance platform, TENS) or a treadmill training group to improve proprioception and balance.

- Results showed significant improvement in proprioception variation and balance in both groups ( $p < 0.05$ ); multi-sensorimotor training showed more significant changes in proprioception ( $p = 0.002$ ) and balance ability ( $p = 0.033$ ).

SENSORY CHANGES AFTER BRAIN INJURY

---

---

---

---

---

---

---

---

42

## VESTIBULAR & PROPRIOCEPTION



### Assessments for balance

- Berg Balance Scale (BBS)
- Sensory Organization Test (SOT)
- Functional Gait Assessment (FGA)
- Dynamic Gait Index (DGI)
- Timed Up and Go (TUG)
- Rivermead Mobility Index
- Functional Reach Test
- Bruininks-Oseretsky Test of Motor Proficiency (Balance subtests)

### Assessments for proprioception

- Active/Passive Joint Position Recognition
- Cervical Joint Position Error Test
- Romberg Test
- Heel-Shin Test
- Finger-Nose-Finger Test
- Thumb Localization Test

SENSORY CHANGES AFTER BRAIN INJURY

---

---

---

---

---

---

---

---

43

## OLFACTORY/GUSTATORY



Screening for olfactory and gustatory dysfunction may involve:

- Asking questions regarding changes in taste, diet, and meals in clinical interview
- Objective quality assessments of smell and taste (University of Pennsylvania Smell Identification Test (UPSIT), Alberta Smell Test (AST))
- Formal testing with taste strips or electrogustometry

Attention to patient nutrition and activity has shown to reduce risk of malnutrition, weight loss, improve certain functional outcomes, and quality of life.

Education to care providers and implementing practices to identify eating difficulties, providing supplementary nutrition to reduce incidence of malnutrition are important in the acute phase to facilitate rehabilitation progress.

SENSORY CHANGES AFTER BRAIN INJURY

(Dutta et al., 2013)

---

---

---

---

---

---

---

---

44

## OLFACTORY/GUSTATORY



Treating olfactory and gustatory dysfunction:

- Steroid (prednisolone) has been shown to have significant effects on recovery rates and may minimize any further decline of olfactory dysfunction and promote regeneration (Proskynitopoulos, et al., 2016).
- Meta analysis of 13 studies' effect on smell training using the Sniffin' Sticks Test. Positive, statistically significant effects of olfactory training with large effects on identification, discrimination, and threshold-discrimination-identification (TDI) score, and small-moderate effect for threshold of odor detection. Significant improvement noted across all olfactory disease subgroups (Parkinson's disease, olfactory disease, post-infectious olfactory loss, patients with post-infectious and post-traumatic olfactory loss, healthy controls) (Sorokowska, et al., 2017).
- May look to COVID-19 research on olfactory and gustatory dysfunction for further understanding of mechanism and treatment options.

SENSORY CHANGES AFTER BRAIN INJURY

---

---

---

---

---

---

---

---

45

## TACTILE



Literature review of post-stroke sensory dysfunction (PSSD) by Gandhi et al. (2021):

Rehabilitation of PSSD is provided by using sensory inputs, categorized as:

1. **Sensory stimulation approach (SSA)** – Passive, externally applied sensory stimulation (electrical, thermal, tactile, repetitive peripheral magnetic stimulation, etc.)
2. **Sensory retraining approach (SRA)** – Active, graded re-education using principles of learning (mirror therapy, graded sensory discriminative activities, mental imagery, graded sensory recognition strategies, combined protocols, etc.)

Of the studies reviewed, improvements reported with mechanical sensation and motor function with alternating hot/cold stimulation, improved somatosensation and motor impairment with intermittent pneumatic compression (IPC), improved quality of life, motor function, tactile and proprioceptive sense with intensive tactile stimulation, functional re-education, and mental imagery over multiple therapy sessions.

SENSORY CHANGES AFTER BRAIN INJURY

---

---

---

---

---

---

---

---

46

## TACTILE



Gironda et al. (2009) on management of chronic pain, post-trauma headache:

- Pharmacological management is complex (multiple injuries, risk of polypharmacy)
  - *Noiceptive: NSAIDs, topical agents, acetaminophen, antispasticity, opioids*
  - *Neuropathic: Topical agents, anticonvulsant, antidepressants, anticonvulsants, opioids*
- Interventional techniques (trigger point injections, nerve blocks, epidural steroids)
- Rehabilitative interventions (thermal modalities, ROM/stretching, US therapy, TENS), graded exercise approach with pain neuroscience education (PNE)
- Cognitive-Behavioral interventions (address emotional/cognitive issues, pain coping skills, address pain-related fear and activity avoidance, medication abuse, PNE)
- Post-trauma headache treatment: NSAIDs, acetaminophen with caffeine or sedative in a single medication, abortive/preventative medications for migraine, relaxation training, biofeedback, trigger identification, stress-management, spinal manipulation, lifestyle regulation, sleep hygiene

SENSORY CHANGES AFTER BRAIN INJURY

---

---

---

---

---

---

---

---

47

## INTEROCEPTION



Research by Desdentado et al. (2023):

Little research that explores changes in ability to regulate emotions following ABI; however, it is known that individuals with TBI are more likely to have maladaptive coping mechanisms, difficulty controlling behavior, and depressive symptomatology.

Interoception and emotional regulation are especially relevant to ABI, since poor insight into deficits ("anosognosia") is a common sequela in brain injury.

Findings lend support to the application of mindfulness-based approaches on emotional processing (meditation, breath work, body scan, etc.).

SENSORY CHANGES AFTER BRAIN INJURY

---

---

---

---

---

---

---

---

48



## INTEROCEPTION



Raimo et al. (2019):

Study on a small sample of patients with unilateral BI that completed the Self-Awareness Questionnaire (SAQ) showed that patients with right brain lesions and extra-personal and/or personal neglect performed significantly worse than controls and patient with left brain lesions on visceral sensations.

This deficit of awareness of visceral sensations possibly reflects differences in brain activation patterns for processing somatic and visceral sensation.

Assessing interoceptive awareness could improve rehabilitation gains and promote greater accuracy of body signals.

SENSORY CHANGES AFTER BRAIN INJURY

---

---

---

---

---

---

---

---

49

### III. EDUCATION OF SENSORY CHANGES FOLLOWING BRAIN INJURY.

"Please take responsibility for the energy you bring into this space."  
- Jill Bolte Taylor

---

---

---

---

---

---

---

---

50



## IMPACT OF SENSORY DYSFUNCTION

Subjective sensory hypersensitivity is known to reduce one's quality of life:

- Social isolation
- Reduced mental health (increased negative affect, depression)
- Reduced physical health (fatigue, sleep disturbance)
- Inability to carry out instrumental activities of daily living (IADL's, leisure)

*"After my stroke, I had to start wearing sunglasses while watching television because of my hypersensitivity to the bright light of the television screen and the flashing images on the screen."*

*"When I am surrounded by a lot of sensory stimuli, I feel tired, nauseous, and anxious."*

*"Everything tastes like dirt"*

*"When I sit in a moving elevator or wheelchair, I feel very uncomfortable. The feeling of those movements is terrible."*

*"Just a small increase in temperature causes me to feel overwhelmed. I do not like when my husband turns on the heating in our house, even when it is cold outside."*

(Thielen et al., 2023;

SENSORY CHANGES AFTER BRAIN INJURY

Dutta et al., 2013)

---

---

---

---

---

---

---

---

51

## QUALITY OF LIFE



Lew et al. (2009): Study of 62 patients with diagnosed sensory impairments (53, **85.5%**, DSI 20, **32.3%**), using the Functional Independence Measure (**FIM**) as the outcome measure:

- At discharge, the DSI group presented with lower scores in all 4 subcomponents of the motor FIM (self-care, mobility, locomotion, sphincter control). Group mean values differed significantly in the self-care subcomponent ( $F(3, 58) = 2.93, p < 0.05$ ).

Shepherd et al. (2020): Health-related quality of life was measured 12 months post-TBI with the Short-Form 36 Health Survey (**SF-36**) and measures of noise, light sensitivity were taken from the Rivermead Post-concussion Symptom Questionnaire (**RPQ**).

- Approximately 1 in 5 individuals are struggling with either noise, light, or both, 12 months following a mTBI. Sensory sensitivities may impede ability to engage in and perform tasks affecting community participation (such as return to meaningful employment).

SENSORY CHANGES AFTER BRAIN INJURY

---

---

---

---

---

---

---

---

52

## QUALITY OF LIFE



Impact of sensory impairments (especially DSI) on neuro-psychological performance:

- Patients with TBI-related, but undetected hearing impairment or DSI may be at risk of being labeled with cognitive or memory impairment.
- Minimal visual blur (20/40 acuity) reduces performance on neuropsych testing (Lew et al., 2010).

Patients report that fatigue plays a role in sensory hypersensitivity (more sensitive to sensory stimuli when tired). They describe physical consequences (headache, fatigue, feeling restless, moist hands, nausea), cognitive consequences (less focus and concentration, feeling distracted), and emotional consequences of sensory hypersensitivity (feeling more irritated, angry, frustrated, acting rude towards others in the environment) (de Sain et al., 2023).

SENSORY CHANGES AFTER BRAIN INJURY

---

---

---

---

---

---

---

---

53



## EDUCATION

Identifying sensory impairment during the acute stage and sub-acute stages of mTBI is important for best-practice rehabilitation and patient quality of life.

- Standardized assessment protocols are lacking; however, we can use subjective report (clinical interview), post-concussion inventories (RPQ), and other tools identified in the literature to address sensory dysfunction.
- Dual sensitivity impairment of noise and light sensitivity should be a focus of clinical assessments of post-concussive symptoms, as these can have a significant influence on long-term recovery and outcome (Shepherd et al., 2020).

Patient education plays a critical role in management of chronic conditions. Research shows that even basic information can improve patient and caregiver self-efficacy and motivation for rehabilitation (Hart et al., 2018).

SENSORY CHANGES AFTER BRAIN INJURY

---

---

---

---

---

---

---

---

54



## ADDITIONAL RESOURCES

Lost & Found: Dealing with Sensory Overload After Brain Injury B. J. Webster, Lash & Assoc. <https://www.brainline.org/article/lost-found-dealing-sensory-overload-after-brain-injury>

Sensory Overload ('Hypersensitivity') after Acquired Brain Injury from the Scottish Acquired Brain Injury Network <https://www.acquiredbraininjury-education.scot.nhs.uk/impact-of-abi/sensory-overload-hypersensitivity/>

Pain after Stroke <https://www.stroke.org.uk/stroke/effects/physical/pain-after-stroke>

The Vestibular Traumatic Brain Injury (TBI) Connection <https://vestibular.org/article/diagnosis-treatment/types-of-vestibular-disorders/tbi/>

Supporting and Enhancing NICU Sensory Experiences <https://chan.usc.edu/nicu/sense>

SENSORY CHANGES AFTER BRAIN INJURY

---

---

---

---

---

---

---

---

55



## READING RECOMMENDATIONS

The Ghost in My Brain: How a Concussion Stole My Life and How the New Science of Brain Plasticity Helped Me Get It Back by Clark E. Elliott

Over My Head: A Doctor's Own Story of Head Injury from the Inside Looking Out by Claudia L. Osborn

My Stroke of Insight: A Brain Scientist's Personal Journey by Jill Bolte Taylor

Volume Control: Hearing in a Deafening World by David Owen

Fixing My Gaze: A Scientist's Journey Into Seeing in Three Dimensions Paperback by Susan R. Barry

The Man Who Mistook His Wife for a Hat and Other Clinical Tales by Oliver Sacks

SENSORY CHANGES AFTER BRAIN INJURY

---

---

---

---

---

---

---

---

56



## REFERENCES

Abusamak, M., & Alrawashdeh, H. M. (2021). Post-Concussion Syndrome Light Sensitivity: A Case Report and Review of the Literature. *Neuro-ophthalmology (Aeolus Press)*, 46(2), 85-90. <https://doi.org/10.1080/01658107.2021.1983612>

Alwis, D. S., & Rajan, R. (2014). Environmental enrichment and the sensory brain: the role of enrichment in remediating brain injury. *Frontiers in systems neuroscience*, 8, 156. <https://doi.org/10.3389/fnys.2014.00156>

Bohnen, N., Twijnstra, A., Wijnen, G., & Jolles, J. (1991). Tolerance for light and sound of patients with persistent post-concussional symptoms 6 months after mild head injury. *Journal of neurology*, 238(6), 443-446. <https://doi.org/10.1007/BF00934653>

Brosseau-Lachaine, O., Gagnon, I., Forget, R., & Faubert, J. (2008). Mild traumatic brain injury induces prolonged visual processing deficits in children. *Brain injury*, 22(9), 657-68. <https://doi.org/10.1080/02699950802203353>

Carrier, S. L., Hicks, A. J., Ponsford, J., & McKay, A. (2021). Managing agitation during early recovery in adults with traumatic brain injury: An international survey. *Annals of physical and rehabilitation medicine*, 64(5), 101532. <https://doi.org/10.1016/j.rehab.2021.101532>

Carron, S. F., Alwis, D. S., & Rajan, R. (2016). Traumatic Brain Injury and Neuronal Functionality Changes in Sensory Cortex. *Frontiers in systems neuroscience*, 10, 47. <https://doi.org/10.3389/fnys.2016.00047>

Chamelian, L., & Feinstein, A. (2004). Outcome after mild to moderate traumatic brain injury: the role of dizziness. *Archives of physical medicine and rehabilitation*, 85(10), 1662-1666. <https://doi.org/10.1016/j.apmr.2004.02.017>

SENSORY CHANGES AFTER BRAIN INJURY

---

---

---

---

---

---

---

---

57



## REFERENCES

- de Sain, A. M., Pelikaan, L. W. M., van Voskuilen, J., Migdis, M., Sommers-Spijkerman, M. P. J., Visser-Meily, J. M. A., & Huenges Wajler, I. M. C. (2023). Sensory hypersensitivity after acquired brain injury: the patient perspective. *Disability and rehabilitation*, 1–8. Advance online publication. <https://doi.org/10.1080/09638288.2023.2251407>
- Desdentado, L., Miragall, M., Llorens, R., Navarro, M. D., & Baños, R. M. (2023). Identifying and regulating emotions after acquired brain injury: the role of interoceptive sensibility. *Frontiers in psychology*, 14, 1268926. <https://doi.org/10.3389/fpsyg.2023.1268926>
- Dutta, T. M., Josiah, A. F., Cronin, C. A., Wittenberg, G. F., & Cole, J. W. (2013). Altered taste and stroke: a case report and literature review. *Topics in stroke rehabilitation*, 20(1), 78–86. <https://doi.org/10.1310/tsr.2013-78>
- Faltynek, P., Marshall, S., Sequeira, K., Teasell, R. (2019). Chapter 6: Motor and Sensory Systems [PDF]. Retrieved from <https://erabi.b-utn.net/wp-content/uploads/2019/09/Chapter-6-Motor-Sensory.pdf>
- Gandhi, D.B.C., Sebastian, I.A., & Bhanot, K. (2021). Rehabilitation of Post Stroke Sensory Dysfunction—A Scoping Review. *Journal of Stroke Medicine*, 4(1), 25–33. doi:[10.1177/2516608520984296](https://doi.org/10.1177/2516608520984296)
- Gironda, R. J., Clark, M. E., Ruff, R. L., Chait, S., Craine, M., Walker, R., & Scholten, J. (2009). Traumatic brain injury, polytrauma, and pain: challenges and treatment strategies for the polytrauma rehabilitation. *Rehabilitation psychology*, 54(3), 247–258. <https://doi.org/10.1037/a0016909>
- Hammerle, M., Swan, A. A., Nelson, J. T., & Treleven, J. M. (2019). Retrospective Review: Effectiveness of Cervical Proprioception Retraining for Dizziness After Mild Traumatic Brain Injury in a Military Population With Abnormal Cervical Proprioception. *Journal of manipulative and physiological therapeutics*, 42(6), 399–406. <https://doi.org/10.1016/j.jmpt.2018.12.002>

SENSORY CHANGES AFTER BRAIN INJURY

58



## REFERENCES

- Hart, T., Driver, S., Sander, A., Pappadis, M., Dams-O'Connor, K., Bocage, C., Hinkens, E., Dahdah, M. N., & Cai, X. (2018). Traumatic brain injury education for adult patients and families: a scoping review. *Brain injury*, 32(11), 1295–1306. <https://doi.org/10.1080/02699952.2018.1493226>
- Hoffman, A. N., Lam, J., Hovda, D. A., Giza, C. C., & Fanelow, M. S. (2019). Sensory sensitivity as a link between concussive traumatic brain injury and PTSD. *Scientific reports*, 9(1), 13841. <https://doi.org/10.1038/s41598-019-50312-w>
- Irvine, K. A., & Clark, J. D. (2018). Chronic Pain After Traumatic Brain Injury: Pathophysiology and Pain Mechanisms. *Pain medicine (Malden, Mass.)*, 19(7), 1315–1333. <https://doi.org/10.1093/pm/pny153>
- Janzen, S., McIntyre, A., Meyer, M., Sequeira, K., & Teasell, R. (2014). The management of agitation among inpatients in a brain injury rehabilitation unit. *Brain injury*, 28(3), 318–322. <https://doi.org/10.3109/02699952.2013.860478>
- Kharrazian D. (2015). Traumatic Brain Injury and the Effect on the Brain-Gut Axis. *Alternative therapies in health and medicine*, 21 Suppl 3, 28–32. <https://www.researchgate.net/publication/281644166>
- Kumar, S., Rao, S. L., Nair, R. G., Pillai, S., Chandramouli, B. A., & Subbakrishna, D. K. (2005). Sensory gating impairment in development of post-concussive symptoms in mild head injury. *Psychiatry and clinical neurosciences*, 59(4), 466–472. <https://doi.org/10.1111/j.1440-1819.2005.01400.x>
- Lee, H., Kim, H., Ahn, M., & You, Y. (2015). Effects of proprioception training with exercise imagery on balance ability of stroke patients. *Journal of physical therapy science*, 27(1), 1–4. <https://doi.org/10.1589/jpts.27.1>

SENSORY CHANGES AFTER BRAIN INJURY

59



## REFERENCES

- Lew, H. L., Garvert, D. W., Pogoda, T. K., Hsu, P. T., Devine, J. M., White, D. K., Myers, P. J., & Goodrich, G. L. (2009). Auditory and visual impairments in patients with blast-related traumatic brain injury: Effect of dual sensory impairment on Functional Independence Measure. *Journal of rehabilitation research and development*, 46(6), 819–826. <https://doi.org/10.1682/jrdd.2008.08.0129>
- Lew, H. L., Jerger, J. F., Guillory, S. B., & Henry, J. A. (2007). Auditory dysfunction in traumatic brain injury. *Journal of rehabilitation research and development*, 44(7), 921–928. <https://doi.org/10.1682/jrdd.2007.09.0140>
- Lew, H. L., Wehling, J., Myers, P. J., Pogoda, T. K., & Goodrich, G. L. (2010). Dual sensory impairment (DSI) in traumatic brain injury (TBI)—An emerging interdisciplinary challenge. *NeuroRehabilitation*, 26(3), 213–222. <https://doi.org/10.3233/NRE-2010-0242>
- Lim C. (2019). Multi-Sensorimotor Training Improves Proprioception and Balance in Subacute Stroke Patients: A Randomized Controlled Pilot Trial. *Frontiers in neurology*, 10, 157. <https://doi.org/10.3389/fneur.2019.00157>
- Multidimensional Assessment of Interoceptive Awareness. (n.d.). UCSF Osher Center for Integrative Medicine. <https://oshcr.ucsf.edu/research/maia>
- Nampiaparampil D. E. (2008). Prevalence of chronic pain after traumatic brain injury: a systematic review. *JAMA*, 300(6), 711–719. <https://doi.org/10.1001/jama.300.6.711>
- Pineda, R., Raney, M., & Smith, J. (2019). Supporting and enhancing NICU sensory experiences (SENSE): Defining developmentally-appropriate sensory exposures for high-risk infants. *Early human development*, 133, 29–35. <https://doi.org/10.1016/j.earhumerdev.2019.04.012>

SENSORY CHANGES AFTER BRAIN INJURY

60



## REFERENCES

Pistola, F., Carolei, A., Sacco, S., Conson, M., Pistorini, C., Cazzulani, B., Stewart, J., Franceschini, M., & Sarà, M. (2015). Contribution of interoceptive information to emotional processing: Evidence from individuals with spinal cord injury. *Journal of neurotrauma*, 32(24), 1981–1986. <https://doi.org/10.1089/neu.2015.3897>

Proskynitopoulos, P. J., Stippler, M., & Kasper, E. M. (2016). Post-traumatic anosmia in patients with mild traumatic brain injury (mTBI): A systematic and illustrated review. *Surgical neurology international*, 7(Suppl 10), S263–S275. <https://doi.org/10.4103/2152-7806.381981>

Preusser, S., Thiel, S. D., Rook, C., Roggenhofer, E., Kosatschek, A., Draganski, B., Blankenburg, F., Driver, J., Villringer, A., & Plegier, B. (2015). The perception of touch and the ventral somatosensory pathway. *Brain - a journal of neurology*, 138(Pt 3), 540–548. <https://doi.org/10.1093/brain/awu370>

Raimo, S., Boccia, M., Di Vita, A., Iona, T., Cropano, M., Ammendolia, A., Colao, R., Iocco, M., Angelillo, V., Guariglia, C., Grossi, D., & Palermo, L. (2020). Interoceptive awareness in focal brain-damaged patients. *Neurological sciences : official journal of the Italian Neurological Society and of the Italian Society of Clinical Neurophysiology*, 41(6), 1627–1631. <https://doi.org/10.1007/s10072-019-04472-z>

Schofield, P. W., Moore, T. M., & Gardner, A. (2014). Traumatic brain injury and olfaction: a systematic review. *Frontiers in neurology*, 5, 5. <https://doi.org/10.3389/fneur.2014.00005>

Shepherd, D., Landon, J., Kalloor, M., Barker-Colla, S., Starkey, N., Jones, K., Ameratunga, S., Theadom, A., & BIONIC Research Group (2020). The association between health-related quality of life and noise or light sensitivity in survivors of a mild traumatic brain injury. *Quality of life research : an international journal of quality of life aspects of treatment, care and rehabilitation*, 29(3), 665–672. <https://doi.org/10.1007/s11136-019-02346-w>

SENSORY CHANGES AFTER BRAIN INJURY

61



## REFERENCES

Sorokowska, A., Drechsler, E., Karwowski, M., & Hummel, T. (2017). Effects of olfactory training: a meta-analysis. *Rhinology*, 55(1), 17–26. <https://doi.org/10.4193/rhin16.195>

Theodoroff, S. M., Papesch, M., Duffield, T., Novak, M., Gallun, F., King, L., Chesnutt, J., Rockwood, R., Palandri, M., & Hullar, T. (2022). Concussion Management Guidelines Neglect Auditory Symptoms. *Clinical journal of sport medicine : official journal of the Canadian Academy of Sport Medicine*, 32(2), 82–85. <https://doi.org/10.1097/JSM.0000000000000874>

Thielen, H., Huenges-Wajer, I. M. C., Tuts, N., Welkenhuyzen, L., Lafosse, C., & Gillebert, C. R. (2024). The Multi-Modal Evaluation of Sensory Sensitivity (MESSY): Assessing a commonly missed symptom of acquired brain injury. *The Clinical neuropsychologist*, 38(2), 377–411. <https://doi.org/10.1080/13854046.2023.2219024>

Thielen, H., Tuts, N., Lafosse, C., & Gillebert, C. R. (2023). The Neuroanatomy of Poststroke Subjective Sensory Hypersensitivity. *Cognitive and behavioral neurology : official journal of the Society for Behavioral and Cognitive Neurology*, 36(2), 68–84. <https://doi.org/10.1097/WNN.0000000000000341>

Wattiez, A. S., Castonguay, W. C., Gaul, O. J., Walte, J. S., Schmidt, C. M., Reis, A. S., Rea, B. J., Sowers, L. P., Cintrón-Pérez, C. J., Vázquez-Rosa, E., Pieper, A. A., & Russo, A. F. (2021). Different forms of traumatic brain injuries cause different tactile hypersensitivity profiles. *Pain*, 162(4), 1163–1175. <https://doi.org/10.1097/J.PAIN.0000000000002103>

Wiert, L., Luauté, J., Stefan, A., Plantier, D., & Hamonet, J. (2016). Non pharmacological treatments for psychological and behavioural disorders following traumatic brain injury (TBI): A systematic literature review and expert opinion leading to recommendations. *Annals of physical and rehabilitation medicine*, 59(1), 31–41. <https://doi.org/10.1016/j.rehab.2015.12.001>

Zasler, N. D., Katz, D. I., Zafonte, R. D., (2013). *Brain Injury Medicine: Principles and Practice*, 2nd Edition. Demos Medical Publishing.

SENSORY CHANGES AFTER BRAIN INJURY

62

## THANK YOU



Christina Ball, OTR/L, CBIS  
[christinatball@gmail.com](mailto:christinatball@gmail.com)

SENSORY CHANGES AFTER BRAIN INJURY

63